

## SOME RESULTS ON FUZZY DIVISOR CORDIAL GRAPHS

Mohammed M. Ali Al-Shamiri<sup>1-3</sup>, S.I. Nada<sup>2</sup>, A.I. Elrokh<sup>2</sup>, and Yasser Elmshtay<sup>1</sup>

<sup>1</sup>Department of Mathematics, Faculty of Sciences and Arts, Mohayel Assir, King Khalid University, Saudi Arabia

<sup>2</sup>Department of Mathematics, Faculty of Sciences, Munofia University, Monofia, Egypt

<sup>3</sup>Department of Mathematics and computer, Faculty of science, Ibb University, Ibb, Yemen

Copyright © 2019 ISSR Journals. This is an open access article distributed under the *Creative Commons Attribution License*, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**ABSTRACT:** A fuzzy divisor cordial labeling of a fuzzy simple graph  $G = (\sigma, \mu)$  be a bijection  $\sigma$  from  $V$  to  $[0,1]$  such that if each edge  $uv$  is assigned the label  $d$  if either  $\sigma(u) \mid \sigma(v)$  or  $\sigma(v) \mid \sigma(u)$  where  $d \in (0,1)$ , and the label 0 otherwise, then the number of edges labeled with 0 and the number of edges labeled with  $d$  differ by at most 1. If a graph has a fuzzy divisor cordial labeling, then it is called fuzzy divisor cordial graph. In this paper, we proved that path, cycle, wheel graph, star graph, some complete bipartite graphs, shell graph  $S_n, S(K_{1,n})$  graph, graph  $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$  and the Helm  $H_n$  graph are fuzzy divisor cordial graph.

**KEYWORDS:** Fuzzy Divisor Cordial Labeling, Fuzzy Divisor Cordial Graph, Fuzzy Labeling.

### 1 INTRODUCTION

The origin of graph labeling problem dates back to the thirteenth century when the Chinese Mathematician Yang Hui and others studied the labeling of geometric figures, which are now classified as plane graphs. The efforts to find solutions to many practical problems in real life situations have also led to the development of several graph labeling methods - graceful, harmonious, prime, divisor, magic, anti-magic, cordial, product cordial, prime cordial etc. Various labeling methods also generate so many new types of graphs. Familiar family graphs have become increasingly useful from mathematical models to a wide range of applications such as design of good radar type codes with automatic correlation properties, identification of ambiguity in X-ray crystallography, formulation of network communication system processing, and optimal circuit layout determination. Diversification of graphic labeling in some references. Marking is a strong connection between number theory and graph structure. By combining the concept of schism in number theory and the concept of a quiet mark in graphing, where the concepts of divisibility play an important role in number theory. Cahit introduced the concept of cordial labeling. And he proved some results related to cordial labeling. Sundaram, Ponraj and Somasundaram in [5] have introduced the notion of prime cordial labeling. And proved that some graphs are prime cordial. Motivated by the concept of prime cordial labeling, R. Varatharajan in [10] introduced a new special type of cordial labeling called divisor cordial labeling. And he proved the standard graphs such as path, cycle, wheel, star and some complete bipartite graphs are divisor cordial. Also show that a complete graph  $K_n$  is not divisor cordial for  $n \geq 7$ . More than that, the special graphs  $G * K_{1,n}$ ,  $G * K_{2,n}$  and  $G * K_{3,n}$  are divisor cordial. M. Sumathi and A.A gwin in [9] charles introduced a new concept called fuzzy divisor cordial labeling. It is a conversion of crisp graph into fuzzy graph under the new condition namely fuzzy divisor cordial labeling, he proved that a complete  $n$ -nary tree is a fuzzy divisor cordial graph, and Every complete graph  $k_n$  is a fuzzy divisor cordial graph. In divisor cordial labeling it is not possible to label all the crisp graphs due to the condition of its definition. Suppose if we consider a graph of size 5, it will be possible to label all the vertices as in the combination of vertex set  $\{1,2,3,4,5\}$ . So for  $n$  vertices, we need to label all the vertices as a combination of all the vertices without repetition, without neglecting any vertex among them. Here discussion about the edge labeling is trivial. So it is clear that all the crisp graphs can't be divisor cordial graphs. However in fuzzy divisor cordial graph for any vertices we can label any fuzzy membership value from  $[0,1]$ . Since the interval consists of infinite number of terms, there are infinite number of chances for labeling a vertex in fuzzy divisor cordial labeling. Fuzzy graphs have been witnessing a tremendous growth and finding applications in many branches of engineering and technology so far. In

[8]Rosenfeld has obtained several concepts in fuzzy graph like fuzzy bridges, fuzzy paths, fuzzy cycles, and fuzzy trees and established some of their properties. In this paper we have introduced some results on fuzzy divisor cordial graph , as an application of labeling fuzzy numbers for the graphs under some new conditions. Fuzzy graph is the generalization of the crisp graph. So it is necessary to know some basic definitions and concepts of fuzzy graph and fuzzy divisor cordial graphs.

## 2 PRELIMINARIES

### DEFINITION 2.1 [4]

A Graph  $G(V, E)$  is of vertices and edges. The vertex set  $V(G)$  is non-empty set and the edge set  $E(G)$  may be empty. A vertex is simply an element of the vertex set and an edge represents a connection between two elements of unordered pair from the vertex set.

### DEFINITION 2.2 [4]

A complete graph  $K_n$  of order  $n$  is a graph in which every two distinct vertices are adjacent

### DEFINITION 2.3 [4]

A graph  $G$  is called bipartite if  $V$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that every edge of  $G$  joins a vertex of  $V_1$  with a vertex of  $V_2$ . If every vertex in  $V_1$  is adjacent to every vertex in  $V_2$ , then  $G$  is said to be complete bipartite graph, denoted by  $K_{m,n}$ , where  $m = |V_1|$  and  $n = |V_2|$ .

### DEFINITION 2.4 [4]

A path of length  $n - 1$ , denoted by  $P_n$ , is a sequence of distinct edges  $v_1v_2, v_2v_3, \dots, v_{n-1}v_n$  with  $v_i v_{i+1} \in E(P_n)$ . A closed path, a path with  $v_1 = v_n$ , is called a cycle.

### DEFINITION 2.5 [4]

A wheel  $W_n$  is defined as  $K_1 + C_{n-1}$ ,  $n \geq 4$ .

### DEFINITION 2.6

The graph  $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$  is the graph obtained by joining apex (central) vertices of stars to new vertex  $x$ .

### DEFINITION 2.7

The subdivision of star  $S(K_{1,n})$  is the graph obtained from  $K_{1,n}$  by attaching a pendent edge to each vertex of  $K_{1,n}$  except root vertex.

### DEFINITION 2.8 [3]

The helm graph  $H_n$  is the graph obtained from an  $n$ -wheel graph by adjoining a pendant edge at each node of the cycle.

### DEFINITION 2.9 [2]

The shell  $S_n$  is the graph obtained by taking  $n - 3$  concurrent chords in cycle  $C_n$ .

### DEFINITION 2.10 [3]

Let  $G = (V, E)$  be a graph. A mapping  $f : V(G) \rightarrow \{0,1\}$  is called binary vertex labeling of  $G$  and  $f(v)$  is called the label of the vertex  $v$  of  $G$  under  $f$ . For an edge  $e = uv$ , the induced edge labeling  $f^* : E(G) \rightarrow \{0,1\}$  is given by  $f^*(e) = |f(u) - f(v)|$ . Let  $v_f(0), v_f(1)$  be the number of vertices of  $G$  having labels 0 and 1 respectively under  $f$  and  $e_f(0), e_f(1)$  be the number of edges having labels 0 and 1 respectively under  $f^*$ .

**DEFINITION 2.11 [1]**

A binary vertex labeling of a graph  $G$  is called a cordial labeling if  $|v_f(0) - v_f(1)| \leq 1$  and  $|e_f(0) - e_f(1)| \leq 1$ . A graph  $G$  is cordial if it admits cordial labeling.

**DEFINITION 2.12 [5]**

A prime cordial labeling of a graph  $G$  with vertex set  $V$  is a bijection  $f$  from  $V$  to  $\{1, 2, \dots, |V|\}$  such that if each edge  $uv$  assigned the label 1 if  $gcd(f(u), f(v)) = 1$  and 0 if  $gcd(f(u), f(v)) > 1$ , then the number of edges labeled with 1 and the number of edges labeled with 0 differ by at most 1.

**DEFINITION 2.13 [10]**

Let  $G = (V, E)$  be a simple graph and  $f : V \rightarrow \{1, 2, \dots, |V|\}$  be a bijection. For each edge  $uv$ , assign the label 1 if either  $f(u) | f(v)$  or  $f(v) | f(u)$  and the label 0 if  $f(u) \nmid f(v)$ .  $f$  is called a divisor cordial labeling if  $|e_f(0) - e_f(1)| \leq 1$ .

If a graph has a divisor cordial labeling, then it is called divisor cordial graph.

**DEFINITION 14 [8]**

A fuzzy graph  $G = (\sigma, \mu)$  is a pair of functions  $\sigma : V \rightarrow [0, 1]$  and  $\mu : V \times V \rightarrow [0, 1]$ , where for all  $u, v \in V$ , we have  $\mu(u, v) \leq \sigma(u) \wedge \sigma(v)$ .

**DEFINITION 2.15 [7]**

A graph  $G = (\sigma, \mu)$  is said to be a fuzzy labeling graph if  $\sigma : V \rightarrow [0, 1]$  and  $\mu : V \times V \rightarrow [0, 1]$  is bijective such that the membership value of edges and vertices are distinct and  $\mu(u, v) \leq \sigma(u) \wedge \sigma(v)$  for all  $u, v \in V$ .

**DEFINITION 2.16 [9]**

Let  $G = (\sigma, \mu)$  be a simple graph and  $\sigma : V \rightarrow [0, 1]$  be a bijection. For each edge  $uv$ , assign the label  $d$  if either  $\sigma(u) | \sigma(v)$  or  $\sigma(v) | \sigma(u)$  and the label 0 if  $\sigma(u) \nmid \sigma(v)$ .  $\sigma$  is called a fuzzy divisor cordial labeling if  $|e_\mu(0) - e_\mu(d)| \leq 1$ , where  $d$  is a very small positive quantity, which is closure to 0 and  $d \in (0, 1)$ .

**3 MAIN RESULTS**

Some result in this section is listed in [3] but we introduced in another labeling and prove also some result are new.

**THEOREM 3.1.**

The path  $P_n$  is a fuzzy divisor cordial.

**PROOF.**

Let  $v_1, v_2, \dots, v_n$  be the vertices of the path  $P_n$ .

We can label these vertices by divided it to many parties.

Part one consists of  $v_i$  vertices, where  $m = 1$ , and  $k = 0, 1, 2, \dots$ , and  $\mu(v_{i_k}) = \frac{(2m-1)2^{k_m}}{10^k}$ ; where  $(2m - 1)2^{k_m} \leq n$  and  $m = 1, k_m \geq 0$ .

Part two consists of  $v_j$  vertices, where  $m = 2$  and  $k = 0, 1, 2, \dots$ ,

and  $\mu(v_{j_k}) = \frac{(2m-1)2^{k_m}}{10^k}$ , where  $(2m - 1)2^{k_m} \leq n$  and  $m = 2, k_m \geq 0$ .

We observe that:  $\frac{(2m-1)2^a}{10^k}$  divides  $\frac{(2m-1)2^b}{10^k}$  ( $a < b$ ) and  $\frac{(2m-1)2^{k_i}}{10^k}$  does not divide  $2m + 1$ .

In the above labeling, we see that the consecutive adjacent vertices have the labels even numbers in numerator of fraction and consecutive adjacent vertices have labels odd and even numbers contribute  $d$  to each edge. Similarly, the consecutive

adjacent vertices have the labels odd numbers in numerator of fraction and consecutive adjacent vertices have labels even and odd numbers contribute 0 to each edge. Thus,  $e_{\mu}(d) = \frac{n}{2}$  and  $e_{\mu}(0) = \frac{n-2}{2}$  if  $n$  is even.

And  $e_{\mu}(d) = e_{\mu}(0) = \frac{n-1}{2}$  if  $n$  is odd.

Hence,  $|e_{\mu}(0) - e_{\mu}(d)| \leq 1$ . Thus,  $P_n$  is a fuzzy divisor cordial.

**THEOREM 3.2.**

The cycle  $C_n$  is fuzzy divisor cordial.

**PROOF.**

Let  $v_1, v_2, \dots, v_n$  be the vertices of the cycle  $C_n$ . We follow the same labeling pattern as in the path, except by interchanging the labels of  $v_1$  and  $v_2$ . Then it follows  $C_n$  is fuzzy divisor cordial.

**THEOREM 3.3.**

The wheel graph  $W_n = K_1 + C_{n-1}$  is fuzzy divisor cordial.

**PROOF.**

Let  $v_1$  be the central vertex and  $v_2, v_3, \dots, v_n$  be the vertices of  $C_{n-1}$ .

Case (i):  $n$  is odd.

Label the vertices  $v_1, v_2, \dots, v_n$  as in the labels of cycle  $C_n$  in the Theorem 2.2, with the same order.

Case (ii):  $n$  is even

Label the vertices  $v_1, v_2, \dots, v_n$  as in the labels of cycle in the Theorem 2.2, with the same order except by interchanging the labels of the vertices  $v_2$  and  $v_3$ .

In both the cases, we see that  $e_{\mu}(0) = e_{\mu}(d) = n - 1$ .

Hence,  $|e_{\mu}(0) - e_{\mu}(d)| = 0$ . Thus,  $W_n$  is fuzzy divisor cordial.

**THEOREM 3.4.**

The star graph  $K_{1,n}$  is fuzzy divisor cordial.

**PROOF.**

Let  $v$  be the central vertex and  $v_1, v_2, \dots, v_n$  be the end vertices of the star  $K_{1,n}$ . Now assign the label  $\frac{2}{10}$  to the vertex, the label  $\frac{1}{10}$  to the vertex  $v_1$  and the vertices  $v_2, \dots, v_n$  label as follows  $e_{\mu}(v_i) = \frac{i+1}{10^i}$ . Then we see that

$$e_{\mu}(0) - e_{\mu}(d) = \begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd} \end{cases}$$

Thus,  $|e_{\mu}(0) - e_{\mu}(d)| \leq 1$  and hence  $K_{1,n}$  is divisor cordial.

**THEOREM 3.5.**

The complete bipartite graph  $K_{2,n}$  is fuzzy divisor cordial.

**PROOF.**

Let  $V = V_1 \cup V_2$  be the bipartition of  $K_{2,n}$  such that  $V_1 = \{x_1, x_2\}$  and  $V_2 = \{y_1, y_2, \dots, y_n\}$ . Now assign the label  $\frac{1}{10}$  to  $x_1$  and  $\frac{p}{10^{\frac{p-1}{2}}}$  where  $p$  the largest prime number of numerator of fraction such that  $p \leq n + 2$  to  $x_2$ , the label  $\frac{i+1}{10^i}$  to the vertices  $y_i, i = 1, 2, \dots, n$ . Then it follows that  $e_{\mu}(0) = e_{\mu}(d) = n$  and hence,  $K_{2,n}$  is fuzzy divisor cordial.

**THEOREM 3.6.**

The complete bipartite graph  $K_{3,n}$  is fuzzy divisor cordial.

**PROOF.**

Let  $V = V_1 \cup V_2$  be the bipartition of  $V$  such that  $V_1 = \{x_1, x_2, x_3\}$  and  $V_2 = \{y_1, y_2, \dots, y_n\}$ . Now define  $\mu(x_1) = \frac{1}{10}, \mu(x_2) = \frac{2}{10}, \mu(x_3) = \frac{p}{10^{\frac{p-1}{2}}}$

Where  $p$  is the largest prime number of numerator of fraction such that  $p \leq n + 3$  and the label  $\frac{i+2}{10^i}$  to the vertices  $y_i, i = 1, 2, \dots, n$ . Then  $e_\mu(0) - e_\mu(d) = \begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd} \end{cases}$

Thus  $K_{3,n}$  is fuzzy divisor cordial.

**THEOREM 3.7.**

$S(K_{1,n})$ , the subdivision of the star  $K_{1,n}$ , is fuzzy divisor cordial.

**PROOF.**

Let  $V(S(K_{1,n})) = \{v, v_i, u_i : 1 \leq i \leq n\}$  and let  $E(S(K_{1,n})) = \{vv_i, v_i u_i : 1 \leq i \leq n\}$ . Label  $v$  by  $\frac{1}{10}$  and define  $\mu$  as  $\mu(v_i) = \frac{2i}{10^i} (1 \leq i \leq n)$  and  $\mu(u_i) = \frac{2i+1}{10^i} (1 \leq i \leq n)$ .

Then we have  $e_\mu(0) = e_\mu(d) = n$ , and therefore  $S(K_{1,n})$  is fuzzy divisor cordial.

**THEOREM 3.8.**

The shell graph  $S_n$  is a fuzzy divisor cordial graph  
For  $n \geq 3$ .

**PROOF:**

Let  $G$  be shell graph  $S_n$ . Let  $v_1, v_2, \dots, v_n$  be vertices of  $G$  with  $v_1$  an apex vertex of  $G$ . Then  $|V(G)| = n$  and  $|E(G)| = 2n - 3$ , label  $U_i$  by  $\mu(v_i) = \frac{2i-1}{10^i} \forall 1 \leq i \leq n$ . Then we have  $e_\mu(0) = n - 2$  and  $e_\mu(d) = n - 1$ .

Therefore  $|e_\mu(0) - e_\mu(d)| \leq 1$ .

Hence,  $S_n$  is fuzzy divisor cordial graph.

**THEOREM 3.9.**

The graph  $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$  is a fuzzy divisor cordial graph.

**PROOF:**

Let  $G$  be a graph  $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$ . Let  $v_1, v_2, \dots, v_n$  be pedant vertices of  $K_{1,n}^{(1)}$  and  $u_1, u_2, \dots, u_n$  be pedant vertices of  $K_{1,n}^{(2)}$ , and  $w$  the a common vertex adjacent to them. Then  $|V(G)| = 2n + 3$ , and  $|E(G)| = 2n + 2$ , label the vertices by  $\mu(u) = \frac{1}{10}, \mu(v) = \frac{p}{10^{\frac{p-1}{2}}}$  where  $p$  the largest prime number from 3 to  $4n + 5$ , and  $\mu(v_i) = \frac{2i+1}{10^i}, \mu(w) = \frac{2n+3}{10^{\frac{p-1}{2}}}$  and  $\mu(u_i) = \frac{2n+2i+3}{10^i}$ . Here we get  $e_\mu(d) = n + 1$  and  $e_\mu(0) = n + 1$

And therefore  $|e_\mu(0) - e_\mu(d)| = 0 < 1$ .

Hence  $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$  is a fuzzy divisor cordial graph.

**THEOREM 3.10.**

The Helm  $H_n$  is a Fuzzy divisor Cordial graph.

**PROOF:**

Let  $G$  be a Helm  $H_n$ . Let  $V$  be the apex vertices of degree 4 and the  $u_1, u_2, \dots, u_n$  be the pendant vertices of  $H_n$ .

Then  $|V(G)| = 2n + 1$ ,  $|E(G)| = 3n$  Vertex labeling  $\mu: V(G) \rightarrow [0,1]$  define as follows:

$$\mu(V) = \frac{1}{10}.$$

Case (i).  $n = 3$

$$f(v_i) = \frac{2i+1}{10^i}, 1 \leq i \leq n \text{ and } f(u_i) = \frac{2i+7}{10^i}, 1 \leq i \leq n.$$

Then  $e_\mu(d) = 4$  and  $e_\mu(0) = 5$ . Therefore,  $|e_\mu(0) - e_\mu(d)| \leq 1$

Case (ii).  $n \geq 4$

Consider  $\lfloor \frac{n}{4} \rfloor = k$  and  $\lfloor \frac{n}{2} \rfloor = m$

$$\mu(v_j) = \frac{2k+2i+1}{10^i}, \mu(u_i) = 3\mu(v_i), 1 \leq i \leq m$$

Labels to the remaining vertices  $v_j, v_{j+1}, \dots, v_n$  where  $j = m + 1, m + 2, \dots, n$  and  $u_j, u_{j+1}, \dots, u_n$

Where  $j = m + 1, m + 2, \dots, n$  by  $\mu(v_j) = \frac{2j+1}{10^j}$  such that  $\mu(v_i) \neq \mu(v_{i+1})$

Where  $m \leq i \leq n - 1$  and  $\mu(v_n) \neq \mu(v_1)$ , and  $\mu(v_i) \neq \mu(u_i) m + 1 \leq i \leq n$ .

Hence we have  $e_\mu(d) = \lfloor \frac{3n}{2} \rfloor$ , and  $e_\mu(0) = \lfloor \frac{3n}{2} \rfloor$ .

Therefore  $|e_\mu(0) - e_\mu(d)| \leq 1$ .

Hence  $H_n$  is fuzzy divisor cordial.

**REFERENCES**

- [1] I. Cahit, Cordial graphs: A weaker version of graceful and harmonious graphs, *Ars combinatoria*, 23, pp. 201-207, 1987.
- [2] Deb P and Limaye N B, On harmonious labeling of some cycle related graphs, *Ars. Combin.*, 65 pp. 177 – 197, 2002.
- [3] J.A.Gallian, A dynamic survey of graph labeling, *The Electronic Journal of Combinatorics*, 22, # DS6, 2017.
- [4] Gross, J. T. and Yellen, J. *Graph Theory and Its Applications*, 2nd ed. Boca Raton, FL: CRC Press, 2006.
- [5] M Sundaram, R Ponraj, S Somasundram "Prime cordial labeling of graphs" *J. Indian Acad. Math*, 27 (2), 373-390, 2005.
- [6] Mordeson J.N., Nair P.S., "Fuzzy Graphs and Fuzzy Hypergraphs" *Physica-Verlag, Heidelberg.*, 2000.
- [7] A.Nagoor Gani and D.Rajalaxmi(a) Subahashini, "Properties of a Fuzzy Labeling Graph", *Applied Mathematical Sciences*, vol. 6, no. 69-27, pp. 3461-3466, 2012.
- [8] A.Rosenfeld, *Fuzzy Graphs*, in: L.A. Zadeh, K.S. Fu, M. Shimura (Eds.), *Fuzzy Sets and Their Applications*, Academic Press, New York, 77-95, 1975.
- [9] M. Sumathi & A. Agwin Charles, "Fuzzy Divisor Cordial Graph" *Inter. J. of Human Resources Management (IJHRM)*, Vol. 5, Issue 6, Oct-Nov, pp. 1-12, 2016.
- [10] R. Varatharajan, S. Navanaeethakrishnan & K. Nagarajan, "Divisor Cordial Graphs, *International J. Math. Combin.* Vol.4, 15-25, 2011.