

## Effect of organic manure and mulch on growth and yield of Plantain (*Musa AAB*) in Ituri Province, in the North-eastern of DR-Congo

Sombo<sup>1</sup>, Ntamwira Bagula Jules<sup>2-3</sup>, Lyadunga Mupenda Kazamba Pascal<sup>3</sup>, and Katunga Musale<sup>1</sup>

<sup>1</sup>Nioka Research Station, National Institute for Agricultural Study and Research, BP 2037 Kinshasa 1, Bunia, Ituri, RD Congo

<sup>2</sup>Mulungu Research Station, National Institute for Agricultural Study and Research, BP 2037 Kinshasa 1, Bukavu, South Kivu, RD Congo

<sup>3</sup>Department of Plant Science, Institute of Agronomic and Veterinary Studies, ISEAV / Mushweshwe, RD Congo

---

Copyright © 2020 ISSR Journals. This is an open access article distributed under the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**ABSTRACT:** The constraints threatening plantain production in the eastern of DR-Congo include soil fertility. An experiment was conducted at the National Institute for Agricultural Study and Research at Nioka Research Station to evaluate the effect of organic manures and mulch on growth and yield parameters of plantain (*Musa AAB*). This study revealed that response of organic manures was significant for all the parameters. Manure application enhanced significantly banana plant growth and yield. The highest bunches weight and yield were produced by the plants receiving manure (24.03t/ha) and mulch (23.24 t/ha) while, lowest bunches and yield were harvested in control plot without fertilizer (17.64 t/ha).

**KEYWORDS:** Banana, Mulch manure, Soil fertility, growth Parameters, yield.

### 1 INTRODUCTION

Bananas and plantains (*Musa* species L. ) are important staple food crops and sources of rural income in most part of sub-Saharan Africa, particularly for the smallholders who grow them in compound gardens. Besides the production in heavily manured compound farms, *Musa* crops are produced in small fields under shifting cultivation and bush fallow, with yields declining rapidly after few production cycles [1]. The all year round fruiting habit of banana puts the crop in a superior position in bridging the 'hunger gap' between annual crop harvests [2]; therefore, contributing significantly to food and income security of people engaged in its production and trade, particularly, in developing countries [3].

Moreover, banana is considered an important food, because of its chemical composition and high content of vitamins and minerals, particularly potassium [3]; [4]; [5]. The pseudostems and leaves are commonly used as mulch in plantations, livestock feed and as wrapping material [5]. Dried leaf bases are extensively used as roofing material for houses, to weave ropes for tethering goats and sheep, and for mattress making [6]. In Eastern and Central Africa, banana is commonly grown by small-scale farmers [7]

Yield decline in plantain cultivation is a common occurrence after few production cycles due to low levels of soil organic matter and increased susceptibility to biotic stress [1]. The constraints threatening plantain production in the eastern of DR-Congo include soil fertility [8].

Better managements of soil fertility are imperative for rehabilitating degraded soil in order to increase crop yields in Africa, especially in South Kivu Province. For example, the strategies of application small doses of fertilizer and planting improved crop varieties were reported [9].

## 2 MATERIAL AND METHODS

An experiment was established Jun 2016 at the Institut National pour l'Etude et la Recherche Agronomiques (INERA) Nioka Research Station, which is located at 02, 211.517 S, 30, 652.322 E and at an altitude of 1653, 6 m above sea level (masl). Nioka Research Station is located in Ituri Province in the North-eastern of DR Congo. This trial was laid out in randomized block design on sandy soil with 3 replications. Plants were spaced at 3 x 2 m. The treatment plot size 6x9m and the replication measured 33x26m. The treatments included three treatments as follows : T1 : Cow manure (3kg/plant); T2 : Mulch (3 kg/plant containing 1.5 kg of *Titonia diversifolia* and 1, 5 kg of *Imperata cylindrica*) and To : control (no fertilizer). 5 kg of farmyard were applied per planting hole for all treatment before planting for both T1 and 2. The manure and mulch were applied after planting and every 6 months (twice before harvest). Each treatment contained 9 banana plants.

Data were collected on the 9 growing plants treatment for recording. Planting, flowering and harvested dates were recorded. Banana growth parameters were assessed in the net plots at flowering stage, the measurements included plant height and pseudostem circumference at soil level. The number of functional banana leaves was also counted at flowering stage. Yield per plot was calculated by multiplying respective mean bunch weight with number of plants (9) in a plot and was expressed in kilograms. Yield per hectare was calculated by multiplying respective mean bunch weight with number of plants per hectare and it was expressed in tons per hectare.

All data were subjected to analysis of variance using the genstat software package [10]. The LSD test was used for means separation. In addition the relation between growth and yield parameters were assessed.

## 3 RESULTS AND DISCUSSION

### 3.1 BANANA GROWTH TRAITS

Organic fertilizers application enhanced plant height, pseudostem circumference at flowering and number of leaves per plant, although this effect was more pronounced for manure than for mulch application (Table 1). The organic fertilizers reduced number of days from planting to flowering in comparison with the control treatment. The result indicates that leaf area influences overall plant growth and confirms observations made by [11]. At banana flowering stage there was no significant effect of manure treatment on banana plant growth in comparison with mulch treatment. The height, circumference of flowering plants and the total number of leaves were lower in control treatment than in the treatment with organic fertilizers, but not significantly effect was observed between the different treatments for circumference and the number of leaves per plant. Contrary, Plants in amended plot with manure were significantly ( $P < 0.05$ ) early flowering (572.5 days) compared to the other treatments. These results corroborates the observation made by [11]; [12]; [13]; [14]; [15]; [16], he revealed that, fertilizer applications significantly affected banana pseudostem height, girth, leaf area and flowering dates.

**Table 1. Effect of organic fertilizers on banana plant height, circumference, Number of days at flowering times and number of produced les per plants**

Treatment	Height (cm)	Circumference (cm)	Number of days at flowering	Number of leaves
Manure	211.9a*	40.3a	572.5b	10.14a
Mulch	190ab	38.8a	638.4a	9.96a
Control	171.1b	34.3a	632.5a	9.82a
LSD	34.64	6.34	29.67	0.9
Fpr	0.069	0.154	0.001	0.785

\*Means in a column followed by the same letter are not significantly different from each other according to LSD at  $P=0.05$ .

### 3.2 BANANA YIELD PARAMETERS

From the data given in the Table 2, it is revealed that, the higher values for yield and yield parameters such as bunch weight, number of hands per bunch, bunches height and yield per hectare (23.64t/ha) were found in treatment which received organic fertilizer (T1 : manure and T2 : mulch), while, minimum was found in control treatment with no fertilizer (T0) which received (17.6 t/ha). The increase in bunch weight could be attributed to increase in yield attributing characters like plant girth, number of functional leaves and leaf area besides higher content of measure nutrients in indexing leaf [15]; [16]; [17], reported that application of NPK at a rate of 600 : 100 : 500 g /mat /yr respectively produced numerically the highest yield. In addition the

organic matter application accelerated bunches maturities (722.1 days) in comparison with mulch (801.6 days) no fertilizer application (777.4 days).

**Table 2. Effect of organic fertilizers on banana yield, bunch weight, bunch length, number of hand and number of days bunches maturity.**

Treatment	Yield (t/ha)	Bunche weight (kg)	Bunche length (cm)	Number of hand	Number of days bunches maturity
Manure	24.03a*	14.1a	56.91a	7.08a	722.1c
Mulch	23.24a	13.94a	58.81a	6.92a	801.6a
Control	17.64b	10.64b	50.1b	6.17b	777.4b
LSD	4	0.98	2.9	0.35	28.57
Fpr	0.005	0.001	0.001	0.001	0.001

\*Means in a column followed by the same letter are not significantly different from each other according to LSD at P=0.05.

#### 4 CONCLUSION

The application of organic fertilizer is one of the important practices for improving banana yield in the poor soil.

From this study it could be concluded that organic fertilization with manure improves the growth and yield parameters.

#### ACKNOWLEDGMENT

Sincere gratitude and appreciation go to the team of persons who helped for data collection of the Institut National pour l'Etude et la Recherche Agronomiques (INERA-NIOKA).

#### REFERENCES

- [1] S. C. Aba, P. K, Baiyeri<sup>1</sup> and A. Tenkouano (2011). Impact of Poultry Manure on Growth Behaviour, Black Sigatoka Disease Response and Yield Attributes of Two Plantain (*Musa* spp. AAB) Genotypes. TROPICULTURA, 2011, 29, 1, 20-27.
- [2] INBAP (1996). Annual Report. INIBAP, Montpellier, France.
- [3] M. Kamira<sup>1</sup>, J. Ntamwira<sup>1</sup>, C. Sivirihauma, W. Ocimati, P. Van Asten, L. Vutseme and G. Blomme (2016). Agronomic performance of local and introduced plantains, dessert, cooking and beer bananas (*Musa* spp. ) across different altitude and soil conditions in eastern Democratic Republic of Congo.
- [4] Ambrin R, Mehrunisa M, Kazi SM, Shamsuddin T, Tanveer AS and Muhammad AK (2017). Nutrient composition of banana fruit as affected by farm manure, composted pressmud and mineral fertilizers. *Pak. J. Bot.* , 49 (1) : 101-108, 2017.
- [5] Karamura EB, (1993). The strategic importance of bananas and plantains in Uganda. In : Gold, C. S. And Gemmil, B (eds). Proceedings of a research co-ordination meeting for biological and integrated control of highland banana and plantain pests and diseases. Cotonou, itacotonou, Benin. Pp 384-397.
- [6] Kamira M, Sivirihauma C, Ntamwira J, Ocimati W, Katungu MG, Bigabwa JB, Vutseme L, Blomme G (2015). Household uses of the banana plant in eastern Democratic Republic of Congo. *J. Appl. Biosci.* 94 : 8915-8929.
- [7] Bagamba F, Senyonga JW, Tushemereirwe WK, Gold CS (1999). Performance and profitability of the banana subsector in Uganda farming systems. In : Proceedings of an International Symposium : Bananas and Food Security, Douala, Cameroon, 10-14 November 1998. INIBAP, Montpellier, France pp. 729-740.
- [8] van Asten PJA, Wairegi LWI, Mukasa D, Uringi NO (2011). Agronomic and economic benefits of coffee–banana intercropping in Uganda’s smallholder farming systems. *Agric. Syst.* 104 (4) : 326-334.
- [9] Ntamwira BJ, Pyame MLD, Kanyenga LA, and Dhed’a DB (2018). The effects of different combinations of herbaceous and shrubs and microdose of fertilizer on bean and maize yields, soil proprieties and carbon sequestration on two degraded soils in the highland of South Kivu, Eastern of DR Congo. *International Journal of Innovation and Applied Studies.* 23 (3), pp. 275-284.
- [10] genstat (2008). Introduction to genstat for Windows. 11th Edition. Lawes Agricultural Trust, Rothamsted Experimental Station, UK.
- [11] Balbín LAG and Zapata JC (2001). Integrated management of black and yellow Sigatoka diseases in plantain cv. Africa. *Infomusa* 10 (2), 3-7.

- [12] Sarathi MW, Jin C, Fransisca CA, Alejandro M, Sean EW and Thomas DS (2015). The relationship between leaf area growth and biomass accumulation in *Arabidopsis thaliana*. *Front Plant Sci*, 6 : 167.
- [13] El-Said ME, Katsuo T, Yositaka S, Asahi O and Yurie U (2016). The Effect of Organic Fertilizers and Flowering Plants on Sheet-Web and Wolf Spider Populations (Araneae : Lycosidae and Linyphiidae) and Its Importance for Pest Control. *J Insect Sci*, 16 (1).
- [14] Khalid A and Rashid A (2009). Effect of NPK fertilizer on growth and yield of banana in Northern Oman. *Journal of Horticulture and Forestry* Vol. 1 (8) pp. 160-167.
- [15] Khair T and Said A (2013). Effects of organic and inorganic fertilizers addition on growth and yield of banana (*Musa* AAA cv. Malindi) on a saline and non-saline soil in Oman. *Journal of Horticulture and Forestry*.
- [16] Ntamwira J, Pypers P, van Asten P, Vanlauwe B, Ruhigwa B, Lepoint P, Dheda B, Monde T and Blomme G (2014). Effect of banana leaf pruning on banana and legume yield under intercropping in farmers' fields in eastern Democratic Republic of Congo. *Journal of Horticulture and Forestry*, vol 6 (9), pp 72-80.
- [17] Aba, P. K. Baiyeri and A. Tenkouano (2011). Impact of Poultry Manure on Growth Behaviour, Black Sigatoka Disease Response and Yield Attributes of Two Plantain (*Musa* spp. AAB) Genotypes. *Tropicicultura*, 29, 1, 20-27.