

Analysis of Physico-Chemical Parameters in Harsi Reservoir Dabra, Gwalior District, Madhya Pradesh

Namrata Pawaiya, D. K. Sharma, and M. K. S. Khushwah

Dept. of Zoology, S.M.S. Govt Model Science College Gwalior, India

Copyright © 2014 ISSR Journals. This is an open access article distributed under the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT: This study was aimed to estimate physico-chemical characteristic of Harsi reservoir. Harsi reservoir located in Dabra, Gwalior district, Madhya Pradesh is constructed on parwati River. Monthly study in physico-chemical parameters such as water temperature, depth, transparency, electrical conductivity, turbidity, total dissolved solids, pH, dissolved oxygen, free carbon dioxide, total alkalinity, total hardness, chlorides, sulphates, nitrate, nitrite, phosphate, silicates, ammonia, BOD, COD, calcium, magnesium, sodium, potassium were analyzed from January 2011 to December 2011. The results indicated that physico-chemical parameters of the water were used for drinking, domestic use, irrigation and pisciculture.

KEYWORDS: Harsi reservoir, physical parameters, chemical parameters, Correlation coefficient.

INTRODUCTION

Quality of an aquatic ecosystem is dependent on the physico-chemical parameters of water. The healthy aquatic ecosystem is dependent on the physico-chemical and biological characteristics (Venkatesharaju et al 2010). Among the fresh water resources of the world pond, river, lakes, reservoirs and wetlands are important because they supply water for the population in the whole year. This they do by collecting rainwater through the water cycle, and sustain the stationary component of the hydrological cycle. They maintain the balance of the ecosystem components on which other systems depend. The lakes and reservoirs represent very complex and fragile ecosystems. The fresh water habitats are considered worthwhile not only as a supply resource but as a living system by which the global ecosystem is balanced. The physico-chemical characteristics play an important role in assessment of the water quality and trophic status of a water body. In Madhya Pradesh, there are many freshwater bodies in the form of rivers, lakes and man-made reservoirs in the state. The small, medium and large reservoirs in this state are estimated to be 1,72,575; 1,67,502 and 1,18,307 ha respectively with a total of 0.458 million ha (Sugunan 1997), contribute to the maximum water spread of all Indian states under manmade lakes. Gwalior and Chambal divisions are rich in water resources and have approximately 54,839 ha water area falling under reservoirs. The determination of good growth in water body include temperature, water temperature, dissolved oxygen, free carbon dioxide, hardness, turbidity, alkalinity etc. other parameters like biological oxygen demand and chemical oxygen demand indicate pollution of a given water body.

STUDY AREA

Harsi reservoir is constructed on Parwati river situated near Harsi village in Dabra, Gwalior District (Madhya Pradesh). It is situated about 100 km from Gwalior. Harsi reservoir was built over the parwati river during 1928-1935. The reservoir is covered by hills and small hillock by its three sides. The reservoir is geographically located between the latitude 25°47' North and longitude to 77°58' East. The maximum height of reservoir is 29.26 m, length is 21,333 m, total catchment area is 777.5 m² and total area cover by reservoir is 308,252 Km³.

METHODS AND MATERIAL

Water Samples were collected from four sampling stations A, B, C, and D from Harsi reservoir. The sampling was done in the first week of every month in early hours of the day i.e, around 9.00 to 11.00 a.m. during January to December 2011. The physico-chemical characteristics of water including water temperature, depth, transparency, pH, dissolved oxygen, free carbon-dioxide, total alkalinity, total hardness, chlorides, calcium and magnesium, were analyzed at the sampling stations, while other parameters including turbidity, electrical conductivity, total dissolved solids, nitrate-nitrogen, nitrite-nitrogen, sulphates, phosphates, silicates, biochemical oxygen demand, chemical oxygen demand, ammonia, sodium and potassium were analyzed in the laboratory within the 6 to 8 hr following the methods of APHA (1995) and Trivedy and Goel (1986). Four sampling stations viz., Station-A, B, C and D were established for analysis of physico-chemical parameters of water covering whole area of reservoir (Fig. 1). The satellite image of the reservoir is in Fig. 2.




Fig 1. Location map and the sampling stations in Harsi reservoir



Fig 2. Satellite image of Harsi reservoir

RESULT AND DISCUSSION

The physico-chemical parameters are determinant of the quality of water. The average value of physico-chemical parameters of water during 2011 are given in table 1. The value of correlation coefficient (r) between physico-chemical parameters of water in harsi reservoir were calculate and given in table 2.

Water Temperature of reservoir water ranged from 18.10 °C to 32.08 °C in different month. temperature increased during summer month and decrease during winter month. Transparency indicates the productive nature of this water on the basis of clarity values as proposed by Sharma and Durve (1991). Transparency of the reservoir ranged from 106.25 cm to 153.50 cm. Electrical conductivity measures the capacity of a substance or solution to conduct electrical current. Electrical conductivity of the reservoir ranged from 177.62 $\mu\text{S}\cdot\text{cm}^{-1}$ to 236.90 $\mu\text{S}\cdot\text{cm}^{-1}$. Turbidity is a measure of water clarity how much the material suspended in water decreases the passage of light through the water. Water turbidity value from 0.91 NTU to 3.14 NTU. The results supported by Dagaonkar and Saksena (1992), Garg et al (2006b) Medudhula.Thirupathaiiah (2012) have also reported high turbidity during rainy season. During rainy season silt, clay and other suspended particles contribute to the turbidity values, while during winter and summer seasons settlement of silt, clay results low turbidity. Total dissolved solids value ranged from 131.25 to 201.0 $\text{mg}\cdot\text{l}^{-1}$ in different month . Similar result have been reported by Rao et al 2003, Kirubavathy et al 2005, Garg et al 2006b.

pH value is very important for plankton growth (Chisty, 2002). During present study water pH values were found 8.19 to 9.09. It is indicating alkalinity nature throughout the study period. This is in accordance with earlier reports by Wetzel (1975), who reported that the value of pH range from 8-9 units in Indian water. Dissolved oxygen is an important aquatic parameter. It plays crucial role in life processes of flora and fauna. In the present study the DO values found from 7.74 to 10.20 $\text{mg}\cdot\text{l}^{-1}$. Range of free carbon dioxide of the reservoir from Nil to 2.76. The carbon dioxide content of water depends upon the water temperature, depth, rate of respiration, decomposition of organic matter, chemical nature of the bottom and geographical features of the terrain surrounding the water body (Sakhare and Joshi, 2002). Total alkalinity is the total sum of carbonates and bicarbonates alkalinity. Total alkalinity range from 30.50 to 76.22 $\text{mg}\cdot\text{l}^{-1}$. In the present study total hardness ranged from 63.0 to 103.0 $\text{mg}\cdot\text{l}^{-1}$ in different seasons. Khanna et al 2010 was also found higher total hardness during monsoon season and lowest total hardness during winter season.

Chloride play an important role in water quality determination. Chloride ranged from 21.43 to 30.86 $\text{mg}\cdot\text{l}^{-1}$. In this study High chloride value during summer season and low value during winter season. Similar observation have also been made by Lendh & Yeragi 2004; and Babar & Raje 2009.

Sulphates is an important constituent of hardness with calcium and magnesium and is one of the key nutrients in the aquatic environment. In the present study sulphates range from 1.11 to 4.16 mg/l⁻¹. The higher value of sulphates recorded in summer month and lower value of sulphates measured in winter month. These results were observed by Kirubavathy et al (2005), Khare et al (2007). During the study Nitrate fluctuated between 0.027 to 0.222 mg/l⁻¹. High concentration of nitrate in drinking water is toxic (Umavathi et al 2007). Nitrogen is a major component of all organic matter. High concentration of nitrates are useful for irrigation but their entry into the water resources increases the growth of nuisance algae, macrophytes and triggers eutrophication and pollution (Trivedy & Goel 1986). High value of nitrate was observed during monsoon season and low during winter season. Similar results were observed by Saxena (1992), Garge et al (2006), Mustapha (2008). Nitrite represents an intermediate form during de-nitrification reactions in the nitrogen cycle. Nitrate concentration was found from 0.075 to 0.290. High concentration of nitrate was found in summer season and low in winter season. Similar results were also observed by Thialaga et al (2005).

Phosphates are one of the most important nutrients and a limiting factor in the maintenance of reservoir fertility. During the study phosphate concentration ranged from 0.004 to 0.032 in different seasons. High concentration of phosphates was observed during monsoon month and low phosphate value during winter. The variation may be due to the various processes like adsorption and desorption of phosphates and buffering action of sediment under varying environmental conditions (Rajasegar, 2003). Silicate is most abundant in sedimentary rocks and therefore, occurs generally in higher concentration in water located in such regions. During the study silicates ranged from 1.76 to 3.95. Higher concentration of ammonia is harmful for fishes and other life. In the present study, the ammonia content varied from 0.39 to 0.74. Higher values of ammonia were found in summer season and lower values in winter season.

Biological Oxygen Demand (BOD) is an important parameter to the oxygen required for degradation of organic matter. During the study period BOD recorded from 0.95 to 3.0 mg/l. Higher values of BOD were found in monsoon season and lower values in winter season. Garg et al., 2010 has also made similar observations in Ramsagar reservoir. Chemical Oxygen Demand is the oxygen consumed by the chemical breakdown of organic and inorganic substances in water. The COD of water increases with increasing concentration of organic matter (Boyd, 1981). In the present study, COD ranged from 18.46 to 36.16 mg/l.

Calcium is an important micronutrient for all organisms. Calcium content ranges from 18.46 to 36.12. The calcium value was high in rainy month and low in winter month. Khanna et al. 2010 has also reported the similar observation. Magnesium is essential for chlorophyll growth and acts as a limiting factor for the growth of phytoplankton (Dagaonkar and Saxena, 1992). Therefore, depletion of magnesium reduces the phytoplankton population. In the present investigation, the magnesium content was found from 3.02 to 7.87 mg/l⁻¹. Sodium is a naturally occurring element in all water bodies. In the present investigation, sodium content was found from 10.02 to 20.79 mg/l. Potassium is also a naturally occurring element, but the concentrations in freshwater bodies remain quite lower than the sodium and calcium. In the present study potassium varied from 1.54 to 4.80 mg/l.

CORRELATION COEFFICIENT BETWEEN PHYSICO-CHEMICAL CHARACTERISTICS

The correlation coefficient (*r*) between physico-chemical parameters are presented in table 2. In the present study show the high degree of positive correlation between temperature and electrical conductivity, temperature and total dissolved solids, temperature and total hardness, temperature and calcium, temperature and magnesium, temperature and chloride, turbidity and total dissolved solids, turbidity and electrical conductivity, turbidity and total hardness, electrical conductivity and total dissolved solids, electrical conductivity and total hardness, electrical conductivity and BOD, electrical conductivity and magnesium, total dissolved solids and total hardness, total dissolved solids and chloride, total dissolved solids and magnesium, COD and chloride, free carbon dioxide and pH, pH and magnesium, total hardness and calcium, total hardness and magnesium, chloride and magnesium, chloride and calcium, calcium and magnesium, BOD and total hardness, COD and chloride. In the present study high degree negative correlation was found between dissolved oxygen and other parameters. Negative correlation between temperature and transparency, temperature and total alkalinity, turbidity and pH, turbidity and total alkalinity, electrical conductivity and pH, electrical conductivity and total alkalinity, total dissolved solids and pH, total alkalinity and calcium, calcium and BOD, calcium and COD, BOD and total alkalinity, COD and Total alkalinity, BOD and COD. Free carbon dioxide has also show negative correlation between most of the parameters. Such a correlation of physico-chemical parameters has also been exhibited by several workers like Jha & Barat (2003), Kumar *et al.* (2007), Garg *et al.*, (2009 & 2010) and Ahangar *et al.* (2012).

Table 1: Average value of physico-chemical parameters of the Harsi reservoir (Jan to Dec 2011)

Parameters	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec
Water temperature (⁰ C)	18.10	22.75	26.0	28.61	30.39	32.08	27.50	26.34	24.43	22.56	21.27	20.48
Transparency (c. m)	141.75	132.25	139.25	127.25	131.50	127.0	114.50	106.25	121.75	138.25	141.75	153.50
E. conductivity ($\mu\text{S.cm}^{-1}$)	183.07	177.62	200.92	209.87	214.55	212.05	234.50	236.90	228.97	204.45	199.65	186.87
Turbidity (NTU)	1.03	1.23	1.66	1.86	2.08	2.27	2.45	2.66	3.14	1.68	1.55	0.91
Total dissolved solids(mgL^{-1})	138.0	143.75	147.50	155.25	162.25	173.25	201.0	194.75	182.50	147.57	146.50	131.25
pH	8.54	8.19	8.35	8.62	8.79	9.09	8.59	8.18	8.19	8.51	8.49	8.55
Dissolved O ₂ (mgL^{-1})	10.20	8.86	8.58	8.38	7.81	7.74	8.33	8.37	8.78	8.86	8.93	9.08
Free CO ₂ (mgL^{-1})	0.897	1.08	0.560	NIL	NIL	NIL	NIL	1.02	2.21	1.40	2.76	1.59
T. Alkalinity (mgL^{-1})	52.45	30.50	39.40	38.45	59.87	76.22	55.77	69.10	45.22	56.0	42.07	61.57
T. Hardness (mgL^{-1})	72.0	74.50	76.50	81.0	85.25	90.50	103.0	94.25	83.0	79.75	70.0	63.0
Chloride (mgL^{-1})	21.43	25.05	29.19	26.83	26.19	30.86	29.49	25.26	21.47	23.12	21.90	21.53
Sulphate (mgL^{-1})	1.68	2.14	2.73	2.66	3.68	4.16	3.06	2.68	2.34	2.33	2.11	1.11
Nitrate-nitrogen (mgL^{-1})	0.027	0.046	0.049	0.071	0.068	0.069	0.113	0.222	0.074	0.057	0.064	0.045
Nitrite-nitrogen (mgL^{-1})	0.081	0.094	0.140	0.098	0.135	0.290	0.212	0.237	0.119	0.075	0.076	0.08
Phosphates (mgL^{-1})	0.007	0.008	0.005	0.010	0.015	0.025	0.032	0.021	0.022	0.014	0.007	0.004
Silicates (mgL^{-1})	1.93	2.47	2.78	3.23	3.46	3.67	3.95	3.45	3.93	2.72	2.22	1.76
Ammonia (mgL^{-1})	0.53	0.51	0.62	0.57	0.66	0.84	0.74	0.59	0.62	0.54	0.48	0.39
BOD (mgL^{-1})	1.39	1.67	1.81	1.98	2.32	2.66	2.79	3.0	2.81	1.27	0.95	0.97
COD (mgL^{-1})	24.62	28.25	34.48	30.34	36.40	44.89	51.08	42.43	41.25	44.36	38.25	21.13
Calcium (mgL^{-1})	18.84	18.46	20.72	30.15	32.72	28.48	36.16	26.49	20.59	22.15	23.48	22.32
Magnesium (mgL^{-1})	3.57	4.80	6.14	5.27	6.42	5.54	7.87	6.58	4.48	3.80	3.98	3.02
Sodium (mgL^{-1})	12.40	13.33	14.78	13.45	13.80	15.54	16.67	17.94	20.79	18.62	14.48	10.02
Potassium (mgL^{-1})	1.54	2.05	2.76	3.23	3.70	4.17	4.53	4.80	3.87	3.40	2.29	2.55

Table 2: Average value of physico-chemical parameters of the Harsi reservoir (January to December 2011)

	WT	TR	EC	TU	TDS	pH	DO ₂	FCO ₂	TA	TH	Cl	SO ₄	NO ₃	NO ₂	PO ₄	Si	NH ₃	BOD	COD	Ca ⁺	Mg ⁺	Na ⁺	K ⁺	
WT	1																							
TR	-0.55	1																						
EC	0.59	-0.83	1																					
TU	0.6	-0.83	0.92	1																				
TDS	0.52	-0.88	0.97	0.9	1																			
pH	0.5	0.17	-0.005	-0.07	-0.14	1																		
DO ₂	-0.93	0.49	-0.58	-0.57	-0.54	-0.40	1																	
FCO ₂	-0.07	0.02	0.25	0.28	0.19	0.15	-0.07	1																
TA	-0.08	0.22	-0.32	-0.37	-0.29	0.29	-0.34	-0.18	1															
TH	0.69	-0.89	0.84	0.77	0.93	0.16	-0.61	-0.20	0.41	1														
Cl	0.81	-0.42	0.34	0.30	0.44	0.450	-0.72	-0.65	0.20	0.64	1													
SO ₄	0.92	-0.53	0.53	0.57	0.57	0.56	-0.84	-0.19	0.38	0.72	0.80	1												
NO ₃	0.33	-0.81	0.75	0.61	0.7	-0.26	-0.38	-0.09	-0.18	0.66	0.20	0.26	-1											
NO ₂	0.62	-0.79	0.71	0.64	0.71	0.11	-0.65	-0.32	0.17	0.78	0.73	0.74	0.8	1										
PO ₄	0.58	-0.8	0.82	0.81	0.84	0.21	-0.53	0.43	0.09	0.92	0.46	0.62	0.55	0.72	1									
Si	0.79	-0.83	0.87	0.92	0.89	0.13	-0.72	0.07	0.26	0.88	0.56	0.76	0.51	0.68	0.86	1								
NH ₃	0.02	0.07	-0.07	0.1	0.01	-0.45	-0.66	-0.07	-0.27	0.80	0.79	0.91	-0.14	-0.18	-0.36	0.81	1							
BOD	0.71	-0.9	0.82	0.88	0.91	0.01	-0.60	-0.06	-0.13	0.87	0.53	0.67	0.65	0.82	0.83	0.91	0.02	1						
COD	0.53	-0.67	0.76	0.76	0.71	0.14	-0.56	0.45	-0.006	0.82	0.45	0.64	0.5	0.6	0.84	0.77	-0.21	0.62	1					
Ca ⁺	0.49	0.28	-0.19	-0.22	-0.34	0.55	-0.69	-0.46	-0.13	0.72	0.59	0.65	0.34	0.16	-0.33	0.63	0.21	-0.12	-0.32	1				
Mg ⁺	0.84	-0.43	0.34	0.31	0.23	0.38	-0.67	-0.42	-0.17	0.85	0.78	0.71	0.22	0.41	0.23	0.74	0.19	0.46	0.26	0.68	1			
Na ⁺	0.24	-0.64	0.72	0.82	0.78	0.29	-0.26	0.52	-0.23	0.58	0.60	0.32	0.74	0.8	0.85	0.68	0.22	0.58	0.79	-0.52	0.4	1		
K ⁺	0.72	-0.8	0.92	0.85	0.84	0.15	-0.74	0.32	-0.05	0.87	0.49	0.64	0.74	0.8	0.85	0.87	-0.11	0.83	0.79	-0.12	0.42	0.64	1	




Fig 3. Showing monthly variations in Water temperature. Fig 4. Showing monthly variations in Turbidity.




Fig 5. Showing monthly variations in Transparency, E. conductivity and Total dissolved solids.




Fig 6. Showing monthly variations in pH and Dissolved oxygen.




Fig 7. Showing monthly variations in Free carbon dioxide




Fig 8. Showing monthly variations in Total alkalinity and Total hardness




Fig 9. Showing monthly variations in Chloride, Calcium and Sodium




Fig 10. Showing monthly variations in Sulphates and Silicate




Fig 11. Showing monthly variations in Potassium and Magnesium.




Fig 12. Showing monthly variations in Phosphate




Fig 13. Showing monthly variations in Nitrate, Nitrite and Ammonia.




Fig 14. Showing monthly variations in BOD




Fig 15. Showing monthly variations in COD

CONCLUSION

In conclusion, All physical and chemical parameters in harsi Reservoir water were within desirable limits. This reservoir have rich amount of nutrients which may be due to agricultural practices being done by farmers in surrounding catchment area of this reservoir and support relatively good biota. The results obtained from the present investigation shall be useful in future management of the reservoir.

ACKNOWLEDGEMENTS

The Authors are grateful to Principal, Head of Department and other staff of Zoology, SMS Govt Modern Science College Gwalior, Madhya Pradesh, India for providing necessary laboratory facilities this research work.

REFERENCES

- [1] APHA (1995). Standard methods for examination of water and wastewater. Nineteenth Edition, Washington D.C.
- [2] Ahangar, I.A., Saksena, D. N., Farooq, M. M. & Ahangar, M.A. (2012). Seasonal variations in physico- chemical characteristics of Anchar lake,
- [3] Kashmir. International Journal of Advance and Biological Research, 2(2), 352-357.
- [4] Babar, H.T. & Raje, G.B. (2009). Water quality indices of some stagnant water bodies in Chiplun, Maharashtra. Journal of Aquatic Biology, 24(2), 124-130.
- [5] Boyd CE (1981). Water quality in warm water fish ponds. Craftmaster Printers Inc., Albama.
- [6] Chisty. N., (2002), Studies on Biodiversity of Freshwater Zooplankton in Relation to Toxicity of selected Heavy Metals. Ph. D. Thesis submitted to M.L Sukhadia Univeristy Udaipur.
- [7] Dagaonkar A, Saksena DN (1992). Physico-chemical and Biological characterization of a temple tank, Kaila Sagar, Gwalior, Madhya Pradesh. J. Hydrobiol. 8(1): 11-19.
- [8] Fokmare AK, Musaddiq M (2002). A study of physico-chemical characteristics of Kapsi lake and Purna River waters in Akola District of Maharastra (India). Nat. Environ. Poll. Tech. 1: 261-263.
- [9] Garg R.K, Rao RJ, & Saksena DN.(2006a).Nutrients and Trophic Status of Ramsagar Reservoir, Datia, Madhya Pradesh Nat. Environ.
- [10] Garg, R. K., Rao, R. J. & Saksena, D.N. (2009). Water quality and conservation management of Ramsagar reservoir, Datia, Madhya Pradesh. Journal of Environmental Biology, 30(5), 909-916.
- [11] Garg, R.K., Rao, R.J., Uchchariya, D., Shukla, G. & Saksena, D.N. (2010). Seasonal variations in water quality and major threats to Ramsagar reservoir, India. African Journal of Environment Science and Technology, 4(2), 61-76.
- [12] Kaushik, S. & Saksena, D.N. (1994). The trophic status and habitat ecology of entomofauna of the water bodies at Gwalior, Madhya Pradesh. In: Perspective in Entomological Research (Agarwal, O.P., ed.), pp. 241-261. Scientific Publishers, Jodhpur.
- [13] Kaushik, S. & Saksena, D.N. (1999). Physico-chemical limnology of certain water bodies of Central India. In: Freshwater ecosystems of India (Vijaykumar, K., ed.), pp. 1-58. Daya Publishing House, New Delhi.
- [14] Khanna, D.R., Kumar, Arun, & Neeraj (2010). Physico-chemical characteristics of a fish pond near Roorkee. International Journal for Enviromental Rehabilitation and Conservation Volume I No. 2 2010 (52-60).
- [15] Khare, S.L., Paul, S.R. & Dubey, A. (2007). A study on water quality of Khomph-Niwari lake at Chhattarpur (M. P.). Nature Environment and Pollution Technology, 6(3), 539-540.
- [16] Kirubavathy, A.K., S. Binukumari, N. Mariamma and T. Rajammal: Assessment of water quality of Orathupalayam reservoir, Erode district, Tamil Nadu. J. Ecophysiol. Occup. Hlth., 5, 53-54 (2005).
- [17] Lendhe, R.S. & Yeragi, S.G. (2004). Physico-chemical parameters and zooplankton diversity of Phirange-Kharbav lake, district Thane, Maharastra. Journal of Aquatic Biology, 19(1), 49-52.
- [18] Medudhula, Thirupathaiiah, Ch.Samatha, Chintha Sammaiah (2012), Analysis of water quality using physico-chemical parameters in lower manair reservoir of Karimnagar district,Andhra Pradesh. International journal of environmental sciences volume 3,no1.
- [19] Mustapha, M.K.(2008).Assessment of the water quality of Oyun reservoir, Offa, Nigeria, using selected physico-chemical parameters Turkis Journal of Fisheries and Aquatic Science, 8, 309-319.
- [20] Ramalingam Manikannan, Subramanian Asokan, Abdul Hameed Mohamed Samsoor Ali, (2012), Seasonal variations of physico- chemical properties of the Great Vedaranyam Swamp, Point Calimere Wildlife Sanctuary, South-east coast of India. African Journal Environmental Science and Technology Vol. 5(9), pp. 673-681, September 2011

- [21] Rajasegar M (2003). Physico-chemical characteristics of the Vellar estuary in relation to shrimp farming. *J. Environ. Biol.*, 24:95-101.
- [22] Rao K D S, Ramakrishniah M, Karthikeyan M, Sukumaran PK (2003). Limnology and fish yield enhancement through stocking in Markonahalli Reservoir (Cauvery River System). *J. Inland Fish. Soc. India*, 35: 20-27.
- [23] Sharma, L. L and Darve, V.S., (1991), Water quality of 26 waters of Rajasthan in relation to phytoplankton. In Proc. Of the second Asian fisheries Forum. Asian fisheries society manila, Philippines, pp 915-918.
- [24] Sakhare, V.B. & Joshi, P.K. (2002). Ecology of Palas-nilegaon reservoir in Osmanabad district Maharashtra. *Journal of Aquatic Biology*, 18(2), 17-22.
- [25] Saxena, Meenakshi (2012), Water quality and trophic status of Raipur reservoir in Gwalior, Madhya Pradesh *Journal of Natural Sciences Research* ISSN 2224-3186 (Paper) ISSN 2225-0921 (Online) Vol.2, No.8, 2012.
- [26] Sugunan, V.V. (1997). Reservoir fisheries in India. FAO Fisheries Technical Paper no, 423.
- [27] Trivedy, R.K. and P.K. Goel: Chemical and Biological Methods for Water Pollution Studies, Environmental Publication, Karad, Maharashtra, India (1986).
- [28] Thilaga, A., S. Subhashini, S. Sobhana and K.L. Kumar: Studies on nutrient content of the Ooty lake with reference to pollution. *Nat. Environ. Poll. Tech.*, 4, 299-302 (2005).
- [29] Uchchariya Dinesh K. (2012), Study of Nutrients and Trophic Status of Tighra Reservoir, Gwalior, *Journal of Natural Sciences Research* ISSN 2224-3186 (Paper) ISSN 2225-0921 (Online) Vol.2, No.8, 2012
- [30] Umavathi, S., Longakumar, K and Subhashini., (2007), Studies on the nutrient content of Sular pond in Coimbatore, TamilNadu, *Journal of ecology and environmental conservation*, 13(5), pp 501-504.
- [31] Venkatesharaju K., Ravikumar.P., Somashekar. R.K., Prakash. K.L., (2010), Physico-chemical and Bacteriological Investigation on the river Cauvery of Kollegal Stretch in Karnataka, *Journal of science Engineering and technology*, 6 (1), pp 50-59.
- [32] Wetzel, R.G. (1975). Primary production and river ecology. Blackwell Scientific Publication, Oxford.