

Systematic Descriptions and Taxonomic studies on Three (3) Species of *Plumeria* in North Central Nigeria

C.U. Aguoru, O.P. Abah, and J.O. Olasan

Department of Biological Sciences, University of Agriculture, Makurdi, Nigeria

Copyright © 2015 ISSR Journals. This is an open access article distributed under the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT: Comparative macro and micromorphology of the leaf and floral features of three species of *Plumeria* was carried out. The aim was to establish the degree of relatedness or separation of the species through the use of macro-morphology, epidermal and floral features as systematic and taxonomic evidences on three species. All the species possessed unvaried anomocytic stomata type on the abaxial surfaces only. Leaf epidermal cells were irregular in shape with the exception of *Plumeria rubra* which possessed pentagonal and hexagonal epidermal cell shapes. Stomata index varied from species to species on the abaxial surfaces and could be applied in their delineation. The index for *Plumeria rubra*, *Plumeria obtusa* and *Plumeria lutea* were 62.34%, 63.31% and 49.57% respectively. Macroscopically, the leaves of all the species were simple and lanceolate, except for *P. obtusa* that is elliptic. The apices of *P. rubra* and *P. lutea* are acuminate while that of *P. obtusa* is acute, the leaf margin of *P. rubra* is undulate while those of *P. obtusa* and *P. lutea* are entire. However, flower type, symmetry, inflorescence, arrangement, calyx colour, corolla shapes, floral formula and floral diagram were unvaried and affirm relatedness and placement in same genus. Variations were however observed in the colour of petals. *P. rubra* is red with yellow centre, while the colour of *P. obtusa* is white with yellow centre. *P. lutea* is yellow fading to white at the tip. The similarities and overlaps observed in the leaf and floral morphology, cell shape, stomata type and stomata index of the species have provided evidence for their genetic and evolutionary relationship and justification for their common grouping at the generic and specific taxa circumscription. These are reported for the first and maiden in its application in three species of *Plumeria*.

KEYWORDS: Micromorphology, *Plumeria*, Stomatal index, Epidermal cells, Taxonomy, Systematics.

1 INTRODUCTION

Genus *Plumeria* belongs to the Apocynaceae family and it is native to the new world. The plants from this genus are widely cultivated in the tropical and subtropical regions throughout the world. *Plumeria* L. is indigenous to tropical America and is found from Southern Mexico to Northern South America and also most abundant in India. The plants are famous for their attractiveness and fragrant flowers. The essential oils from the flowers are used for perfumery (Shaïda *et al.*, 2008).

According to Tung *et al.* (1999) and Scot (2009), *Plumeria* species grow as small ornamental trees in parks, residential and commercial landscapes. *Plumeria* cultivars also adorn roadways and public lands. The decoction of the bark and roots of *P. rubra* is traditionally used to treat asthma, constipation, promote menstruation and reduce fever. The latex is used to soothe irritation (Wiert, 2002). The fruit is reported to be eaten in West Indies. In India, however, it has been used as an abortifacient (Bobbarala *et al.*, 2009). The flowers are aromatic and widely used in pectoral syrups. The flowers decoction of *P. rubra* was reported to use in Mexico for control of diabetes mellitus. The leaves of are used in ulcers, leprosy, inflammations and rubifecient (Kardono *et al.*, 1990). The decoction of the bark of *Plumeria obtusa* is given in varying doses as a purgative or as a remedy against oedemas (Shinde *et al.*, 2014).

Plumeria is generally a small tree growing to about 30 ft high. Its broad, usually round-headed canopy is often about as wide as the tree is tall. The species and hybrids vary somewhat in tree size, compactness, and branching character, leaf and flower size and colour, and deciduousness. The leaves are usually glossy green but may be dull green; they are generally

ovate, may be blunt-tipped (*P. obtusa*) or pointed (*P. rubra* var. *acuminata* or var. *acutifolia*), and range from 2 to 4 inches wide and 8 to 12 inches long. In deciduous types, the leaves fall during wintertime, and new leaves emerge during or following the spring flowering period. *P. obtusa* and its hybrids tend to retain their foliage year-round. The flowers are tubular, expanding into a “pinwheel” of five petals that averages 2–3 inches diameter and may be white, red, yellow, pink, or multiple colours. Flowers of most cultivars are highly fragrant and bloom from March to October. The hybrids differ in their profusion of blooms, with some producing more than 200 flowers per cluster and others only 50–60 flowers. *Plumerias* only occasionally produce seed. When pollinated, the flower produces two hard, narrow, pointed pods up to 7 inches long containing 20–60 winged seeds. Maturation of the seed pods is usually in early spring from a previous season’s pollination (Richard, 1998).

Many studies have been carried out on *Plumeria* species but did not include the leaf and petal epidermal studies. In the findings of Kalam *et al.* (2013) the dried flowers of *Plumeria rubra* f. *rubra* and *Plumeria rubra* f. *lutea* were subjected to successive solvent extraction by soxhlation using n-hexane, chloroform, ethyl acetate and methanol. The extracts were subjected to preliminary phytochemical screening using standard procedures and the data obtained from the flowers of both the species was comparatively evaluated. As essential as genus *Plumeria* is, members of its species must be easily recognized without any difficulty, especially for ornamental or medicinal purposes. Members of this genus are usually identified or recognized with their flowers which are seasonal in appearance. There is therefore, the need for the use of other characteristics which are available all year round, that is, the leaves. Based on the ethno botanical uses of the species, detailed micro and macro morphological studies were carried out. The aim was to establish the phylogenetic relationship among the species (*Plumera rubra*, *Pluneria obtusa*, and *Plumeria lutea*) on the basis of leaf and floral similarities and differences.

2 MATERIALS AND METHOD

Fresh matured leaves and flowers of three species of *Plumeria* were collected from different parts of North Central zone of Nigeria and identified by plant taxonomists in the Botany Unit of the Federal University of Agriculture, Makurdi, Benue State. Each species was phenotypically observed, measured and compared accordingly. Microscopic preparations of the leaf and petal were made to elucidate the epidermal features following standard methods. All preparations were viewed using appropriate objective lenses of the Olympus Microscope. Ten slides were prepared from the leaf and petal of each species. From each slide, ten fields of views were recorded with respect to the stomata type, number of stomata, and number of epidermal cells, shape of epidermal cells and presence or absence of trichome. Mean values of numeric were computed. Description of stomatal complex type was done in accordance with Dilcher (1974). Photomicrographs were taken using an installed digital camera on the microscope. Stomata indices were calculated in percentages using the formula:

$$SI = S/(S + E) \times 100$$

SI = stomata index

S = number of stomata per square millimetre

E = number of epidermal cells per square millimeter

3 RESULTS AND DISCUSSION

The leaves of the species studied were hypostomatic, that is, stomata were present on the abaxial surface but absent on the adaxial surface. The stomata type present was homogenously anomocytic (stomata with five or more subsidiary cells surrounding the guard cells that are not distinguishable from other epidermal cells) (table 1; plate 1, 3, 5). Trichome was absent.

This observation agrees with the work of Chandra *et al.* (1969) who observed similar features in the leaf epidermal studies of *Plumeria rubra*, *Holarrhena pubescens*, *Plumeria alba*, *Wrightia tinctoria*, and *Wrightia tomentosa*. The epidermal cells on all the abaxial and adaxial surfaces were irregular in all the species except in *P. rubra* which possessed pentagonal epidermal cell shape on its abaxial surface and hexagonal epidermal cell shape on the adaxial surface (plate 2). The variation recorded in the epidermal cell aligns with the report of Ghazalah *et al.* (2010) who studied the epidermal cells of 15 species of *Persicaria* Mill. and observed variation among them.

Stomata index varied from species to species on the abaxial surface. The highest stomata index was found on the surface of *P. obtusa* with an index of 63.31% and lowest on the abaxial surface of *P. lutea* with an index of 49.57% (table 1). This

shows that stomata occupied larger proportion of leaf surface in *P. obtusa* and smaller proportion in *P. lutea*. The stomatal index, which indicates the proportion of stomata relative to leaf surface, is also a reliable taxonomic character. This is because it is independent of the changes in epidermal cell size brought about by environmental factors (Metcalf and Chalk, 1988). Similar approach was used by Aguoru and Okoli (2012) to distinguish West Africa species of *Momordica* L using stem and petiole anatomical evidence.

The anatomical features especially the plant epidermis is mildly influenced by environmental conditions and is of high structural diversity. This character represented genetic variations and have been used to solve taxonomic problems in certain other plant groups by taxonomists (Oladele, 1983; Ogunkunle, 2013; Abdulrahman *et al.*, 2009). Their proven genetic stability and high structural diversity have been the bases for their use in identification of many groups (Ogunkunle and Oladele, 2000; Abubakar and Yanusa, 1998).

The three species therefore shared common attributes as revealed in the leaf and floral morphology. Similarity is particularly pronounced in their simple leaf type, pinnately netted venation type, alternate leaf arrangement, cuneate leaf bases, green leaf colours. Similar floral attributes include; bisexuality, cymose flower type, zygomorphic symmetry, green calyx colour, elliptic corolla shapes, floral formula and floral diagram (fig 6). The petals had no stomata on both the abaxial and adaxial surfaces but it was characterized by irregular epidermal cells. However, the differences observed cannot be overlooked. Leaf shapes of *P. rubra* and *P. lutea* were lanceolate while that of *P. obtusa* was elliptic. The leaf apex of *P. rubra* and *P. lutea* were acuminate while that of *P. obtusa* was acute. The leaf margin of *P. rubra* is undulate while those of *P. obtusa* and *P. lutea* were entire (fig 2, 3, 4). The corolla of *P. rubra* was red with yellow centre but that of *P. obtusa* was white with yellow centre. *P. lutea* had yellow corolla fading to white at the tip as (table 2; fig 2, 3, 4).

In conclusion, based on these findings, species of the genus *Plumeria* can readily be distinguished from one another as certain features are considered diagnostic. The overlapping similarities as well as distinguishing characteristics observed among the species have provided evidences for their evolutionary relationship. This work has also provided a justification for their common grouping as well as their divergence at the generic and specific circumscription. The combination of microscopic and macroscopic evidences to establish the phylogenetic relationships among the three species of *Plumeria* has yielded a good result. This may further be enhanced by other approaches such as chemosystematics and molecular characterization.

Table1: Leaf epidermal features

| Species | Leaf surface | Shape of epidermal cells | Stomata type | Stomata index (%) | Trichomes | Leaf length (cm) | Leaf breadth (cm) |
|------------------|--------------------|--------------------------|--------------------------|-------------------|-----------|------------------|-------------------|
| <i>P. rubra</i> | Abaxial Adaxial | Pentagonal Hexagonal | Anomocytic Anomocytic | 62.34 | Absent | 40.0 | 11.8 |
| <i>P. obtusa</i> | Abaxial Adaxial | Irregular Irregular | Anomocytic Anomocytic | 63.31 | Absent | 25.2 | 10.3 |
| <i>P. lutea</i> | Abaxial Adaxial | Irregular Irregular | Anomocytic Anomocytic | 49.57 | Absent | 37.0 | 12.1 |

Table 2: Floral features

| Characters | <i>P. rubra</i> | <i>P. obtusa</i> | <i>P. lutea</i> |
|--------------------|------------------------------|--------------------------------|--------------------------------------|
| Inflorescence | Compound cyme | Compound cyme | Compound cyme |
| Floral symmetry | Zygomorphic | Zygomorphic | Zygomorphic |
| Calyx colour | Green | Green | Green |
| Corolla colour | Red with small yellow centre | White with small yellow centre | Yellow fading to white at the margin |
| Corolla free/fused | Fused | Fused | Fused |
| Nature of sepal | Synsepalous | Synsepalous | Synsepalous |
| Nature of ovary | 2 fused capels to form 1 | 2 fused capels to form 1 | 2 fused capels to form 1 |
| Ovary type | Superior | Superior | Superior |
| Nature of stamen | Epipetalous | Epipetalous | Epipetalous |
| Sex | Bisexual | Bisexual | Bisexual |
| Number of sepals | 5 | 5 | 5 |
| Number of petals | 5 | 5 | 5 |
| Number of anther | 5 | 5 | 5 |
| Petal length (cm) | 4.5 | 4.2 | 6.4 |
| Petal breadth (cm) | 2.2 | 2.1 | 2.7 |




Figure 1: Stomata indices of three species of *Plumeria*




Plate 1




Plate 2




Plate 3




Plate 4




Plate 5




Plate 6




Plate 7




Plate 8




Plate 9




Plate 10

LEGEND:

- Plate 1: *Plumeria rubra* leaf abaxial surface showing anomocytic stomata type and pentagonal epidermal cells
Plate 2: *Plumeria rubra* leaf adaxial surface showing hexagonal epidermal cells without stomata.
Plate 3: *Plumeria obtusa* leaf abaxial surface showing anomocytic stomata type and irregular epidermal cells
Plate 4: *Plumeria obtusa* leaf adaxial surface showing irregular epidermal cells without stomata
Plate 5: *Plumeria lutea* leaf abaxial surface showing anomocytic stomata type and irregular epidermal cells
Plate 6: *Plumeria lutea* leaf adaxial surface showing irregular epidermal cells without stomata
Plate 7: *Plumeria rubra* petal abaxial surface showing irregular epidermal cells without stomata
Plate 8: *Plumeria rubra* petal adaxial surface showing irregular epidermal cells without stomata
Plate 9: *Plumeria obtusa* petal abaxial surface showing irregular epidermal cells without stomata
Plate 10: *Plumeria lutea* petal abaxial surface showing irregular epidermal cells without stomata.



Fig2: *P. rubra* (red petal)



Fig3: *P. obtusa* (white petal)



Fig4: *lutea* (yellow petal)

$$+ \text{♀} \overline{K_{(5)} C_{(5)} A_5 G_{(2)}}$$

Fig.5: Floral formula of *Plumeria* species

LEGEND:

+ = Zygomorphic symmetry; K=Calyx; C=Corolla; A=Androecium; G=Gymnoecium with superior ovary.



Fig 6: Floral diagram of *Plumeria* species

REFERENCES

- [1] Abdulrahman, A.A, Egbedo, F.O, and Oladele, F.A. (2009). Stomatal Complex Types, Stomatal Density, and the Stomatal Index in Some Species of *Dioscorea*. *Niger. J. Bot.* 61(4): 847-850.
- [2] Abubakar, B.Y. and Yanusa, (1998). Epidermal structure and stomatal ontogeny of some species of *Acacia* (*Leguminosae Mimosoidieae*). *Nigeria. J. Bot.*, 11: 117-123.
- [3] Aguru, C.U. and Okoli, B.E. (2012). Comparative stem and petiole anatomy of West African species of *Momordica* L (*Cucurbitaceae*). *African Journal of Plant Science*, 6(15):403-409. DOI:10.5897/AJPS11.309
- [4] Andrew, K.K., Gitta, H. and Drew, M., (2013). *Plumeria*, Propagation from Cutting. Environmental Horticulture Department. ENH1228: 1-2
- [5] Anjum, P., Rubina, A. and Rabab, F., (2007). Stomata types of some Dicots within flora of Karachi, Pakistan, Department of Botany, University of Karachi, Pakistan. Pg (1017).
- [6] Ashraf, M.D., Farooque M. A., Shambhawe, S. and Mazumder, R., (2012). Review on *Plumeria acuminata*, *International Journal of Research in Pharmacy and Chemistry*, 2:2.
- [7] Bobbarala, V., Katikala, P.K., Naidu, K.C, Penmajji, S., (2009). Antifungal activity of selected plant extracts against phytopathogenic fungi *Aspergillus niger* F2723. *Indian Journal of Science and Technology*, 2(4): 87–90.
- [8] Könemann, (2004). The Illustrated AZ of over 10000 garden plants and how to cultivate them. *Botanica*, p 691.
- [9] Bunch, A., (2010). The Exotic *Plumeria*, The World's Number One *Plumeria* Catalog, P 1
- [10] Carol, C. B., (2006). *Plumeria rubra* f. *lutea*. American Native Plants, p 1
- [11] Chandra, V., Kapoor, S.L., Sharma, P.C. and Kapoor, L.D., (1969). Epidermal and venation studies in *Apocynaceae*-1. *Bull. Bot. Sur. India*, 11:286-287.
- [12] Dilcher, D.L., (1974). Approaches to the Identification of Angiosperm Leaf Remains. *Bot. rev.* 40 (1): 1-157.
- [13] Dutta, A.C., (2010). Botany for Degree Student: 6th ed. Oxford University Press. Pg. 25-45, 55-56, 169.
- [14] Ghazalah Y., Mir, A. K., Nighat, S. and Muhammad, Q. H., (2010). Taxonomic Significance of Leaf epidermal Anatomy of Selected *Persicaria* Mill. Species of family Polygonaceae from Pakistan. *African Journal of Biotechnology*, 9(25): 3759-3768.
- [15] Kalam, S., Yegnambhatla, R., Periyasamy, G., Meesa, M., Guggulothu, H. and Yegnambhatla, A., (2013). Comparative Phytochemical and Pharmacological Evaluation of flowers of *Plumeria rubra* L. f. *rubra* and *Plumeria rubra* f. *lutea*. *British Biomedical Bulletin*, pg. 1
- [16] Kapoor, S.L., Sharma, P.C., Chandra, V. and Kapoor, L.D., (1969). Epidermal and venation studies in *Apocynaceae*-II. *Bull. Bot. Surv. India*, 11 (3 and 4): 372-376.
- [17] Kardono, L.B., Tsauri, S., Padmawinata, K., Pezzuto, J.M. and Kinghorn, A.D., (1990). Cytotoxic constituents of the bark of *Plumeria rubra* collected in Indonesia. *Journal of Natural Product*. 53: 1447-1455.
- [18] Kumar, A., Chanda, I. and Singh, A.K., (2009). Extraction and evaluation of pharmacological activity of Saponin extract of *plumeria rubra* leaves. *Pharmacologyonline*, Dept of Pharm. Sci., S.B.S. (PG) IBSR, Dehradun, Uttarakhand, India. 1: 969-974.
- [19] Lawren, S. and Christine, S., (2013). Leaf Venation, structure, function, development, evolution, ecology and applications in the past, present, and future. Department of Ecology and Evolution, University of California Los Angeles, Pg (993-1000).
- [20] Madhavan, V., Mohamed, S.U., Gurudeva, M.R. and Yoganarasimhan, S.N., (2010). *Indian Journal of Natural Products and Resources*, 1(1):38-43.
- [21] Mery, D., Inna, G., Naama, M., Mariana, O.V., Moshe, S., Eran, P., Dani, Z., Efraim, L., Zach, A., David, W., Alexander, V., (2005). Flower Proteome: Changes in protein spectrum during the advanced stages of rose petal development. Faculty of Agriculture, Food and Environmental Quality Sciences, Hebrew University of Jerusalem, Israel, pg 37-45.
- [22] Metcalfe, C.R. and Chalk, L., (1988). Anatomy of the Dicotyledons, 2nd ed. Clarendon Press, Oxford, 1:100-106.
- [23] Ogunkunle, A.T.J. and Oladele, F.A., (2000). Diagnostic Value of Trichomes in Some Nigeria Species of *Ocimum* L., *Hyptis jaccg*, and *Tinnea kotschy* and *Peyr.* (*Lamiaceae*). *J. Applied Sci.*, 3: 1163-1180.
- [24] Ogunkunle, A.T.J., (2013). The Value of Leaf Epidermal Characters in Diagnosing Some Nigeria Species of *Ficus* L. (*Moraceae*). *Research Journal of Botany*, 8: 1-14.
- [25] Oladele, F.A., (1983). Patterns of Cuticular Sculpture in the Hybrid *Cyprinosypris leylandii* (Jackson and Dallimore) Dallimore and its Putative Parents. *New Phytol.*, 94: 293-295.
- [26] Ozoker, E. and Misirh, A., (2011). Research on leaf properties and stomata distribution of some *Pistacia spp*; Department of Horticulture, Faculty of Agriculture Ege University, Turkey, pg 1.
- [27] Peter, D., Jon, A., Christer, B., Piritta, H. and Katri, K., (2008). Leaf Trichome formation and plant resistance to herbivory; marine science Institute, University of California Barbara. Pg 89.

- [28] Richard, A. (1998). Ornamental flowers: Plumeria. Cooperative Extension Service, C/T/A/H/R. Department of Horticulture, College of Tropical Agriculture & Human Resources. OF-24:1-2.
- [29] Richard, A.C., (2005). Ornamentals and flowers: Plumeria. Cooperative Extension Service, C/T/A/H/R. Department of Horticulture, College of Tropical Agriculture & Human Resources, University of Hawai'i at Manoa. OF-32:1-29.
- [30] Scot, N., (2009). Plant disease, Department of Plant and Environment Protection Science. College of Tropical Agriculture and Human Resources University of Hawai'i at Manoa. Pg.1.
- [31] Shaida, F., Salmy, S., Tan, L., Tengku, S. and Tengku, M., (2008). Chemical components of the essential oils from three species of malaysian *Plumeria*. and their effects on the growth of selected microorganisms. *Journal of Bioscience*, 19(2): 1–7.
- [32] Sharma, G.K., (1972). Environmental modifications of leaf epidermis and morphological features in *Verbena canasensis*. *The South-Western Naturalist*, 17(3):221-228.
- [33] Shinde, P.R., Patil, P.S. and Bairagi, V.A.,(2014). Phytopharmacological Review of *Plumeria species*, *Scholars Academic Journal of Pharmacy*, Department of Pharmacognosy, K.B.H.S.S. Trust's Institute of Pharmacy, Malegaon, Dist: Nasik-423105. Maharashtra, India. 3(2): 217-222
- [34] The Southern California Plumeria Society (2011). Caring for Your New Plumeria. Rev. 2, pg 1.
- [35] Tung, A.V., (1999). Pepohon Pinggir Jalan Daun Lebar. Amiza Publishing, Selangor: pg 6–11.
- [36] Wiart, C. (2002). Medicinal Plants of Southeast Asia. Kuala Lumpur, Pearson Malaysia, pg 6-10.